

SECTION 4.10 ANTIDERIVATIVES

- (families of) antiderivatives
- definite integrals
- initial value problems

1. Find the (family of) antiderivatives for the following.

$$(a) f(x) = x^3$$

$$(b) f(x) = 5 \sin(x)$$

$$(c) f(x) = \frac{e^x}{4}$$

$$F(x) = \frac{1}{4}x^4 + C$$

$$F(x) = -5 \cos(x) + C$$

$$F(x) = \frac{1}{4}e^x + C$$

check:

$$F'(x) = \frac{1}{4} \cdot 4x^3 + 0 \\ = x^3 \quad \checkmark$$

$$F' = -5(-\sin(x)) \\ = 5 \sin(x) \quad \checkmark$$

$$F'(x) = \frac{1}{4}e^x \quad \checkmark$$

$$(d) f(x) = \sqrt{2}$$

$$(e) f(x) = \frac{1}{x}$$

$$(f) f(x) = 1 - x + e^x$$

$$F(x) = \sqrt{2}x$$

$$F(x) = \ln|x|$$

$$F(x) = x - \frac{1}{2}x^2 + e^x$$

* Where do absolute value bars come from?

* What principle is being used here?

2. Confirm that $F(x) = \sin^2(2x)$ is an antiderivative of $f(x) = 4 \sin(2x) \cos(2x)$.

$$F'(x) = 2(\sin(2x))'(\cos(2x)) \cdot 2 = 4 \sin(2x)\cos(2x) \quad \checkmark$$

why?

Function	Antiderivative
x^k ($k \neq -1$)	$x^{k+1}/(k+1)$
x^{-1}	$\ln x $
1	x
$\sin(x)$	$-\cos(x)$
$\cos(x)$	$\sin(x)$

Function	Antiderivative
e^x	e^x
$1/(1+x^2)$	$\tan^{-1}(x)$
$\sec^2(x)$	$\tan(x)$
$\sec(x)\tan(x)$	$\sec(x)$
$1/\sqrt{1-x^2}$	$\sin^{-1}(x)$

3. Evaluate the integrals.

$$(a) \int (e^{-x} + \sec^2(x)) dx = -e^{-x} + \tan(x) + C$$

$$(b) \int \frac{x^2 + x^{1/2} + 1}{x^{1/2}} dx = \int \left(x^{3/2} + 1 + x^{-1/2} \right) dx = \frac{2}{5} x^{5/2} + x + 2x^{1/2} + C$$

$$(c) \int \left(\frac{1}{4}x^4 + \sec(x/\pi) \tan(x/\pi) \right) dx = \frac{1}{20} x^5 + \pi \sec\left(\frac{x}{\pi}\right) + C$$

4. Solve the initial value problem if $f'(x) = x + e^x$ and $f(0) = 4$.

$$f(x) = \int (x + e^x) dx = \frac{1}{2} x^2 + e^x + C$$

$$f(0) = \frac{1}{2}(0^2) + e^0 + C = 1 + C = 4. \text{ So } C = 4.$$

$$f(x) = \frac{1}{2} x^2 + e^x + 4$$

5. A particle moving along the x -axis has acceleration $a(t) = 10 \sin(t)$ measured in cm/s^2 . Assume the particle has initial velocity $v(0) = 0$ and initial position $s(0) = 0$, find a function that models its velocity, $v(t)$, and its position $s(t)$.

$$a(t) = 10 \sin(t)$$

$$v(t) = \int a(t) dt = \int 10 \sin(t) dt = -10 \cos(t) + C$$

$$v(0) = -10 \cos(0) + C = -10 + C. \text{ So } C = 10.$$

$$v(t) = -10 \cos(t) + 10$$

$$s(t) = \int v(t) dt = \int (-10 \cos(t) + 10) dt = -10 \sin(t) + 10t + C$$

$$s(0) = -10 \sin(0) + 0 + C = 0. \text{ So } C = 0$$

$$s(t) = -10 \sin(t) + 10$$