

## SECTION 4.10 ANTIDERIVATIVES

**Topics:** (families of) antiderivatives, indefinite integrals, initial value problems

1. Find the (family of) antiderivatives for the following.

(a)  $f(x) = 4x^3$

(b)  $f(x) = 5 \sin(x)$

(c)  $f(x) = \frac{e^x}{4}$

(d)  $f(x) = \sqrt{2}$

(e)  $f(x) = \frac{1}{x}$

(f)  $f(x) = 1 - x + e^x$

2. Is  $F(x) = x + xe^x$  is an antiderivative of  $f(x) = (x + 1)e^x + 1$ ? Show your answer is correct.

Function	Antiderivative
$x^k$ ( $k \neq -1$ )	
$x^{-1}$ for all $x$	
1	
$\sin(x)$	
$\cos(x)$	

Function	Antiderivative
$e^x$	
$1/(1 + x^2)$	
$\sec^2(x)$	
$\sec(x) \tan(x)$	
$1/\sqrt{1 - x^2}$	

3. Evaluate the integrals.

(a)  $\int (x^{1/2} + x^{-7/4}) dx$

(b)  $\int (8e^x + \sec^2(x)) dx$

(c)  $\int \frac{x^2 + x^{1/2} + 1}{x^{1/2}} dx$

4. Is the equality in the box true or false? Explain.

$$\int x \sec^2(x^2 + 1) dx = \tan(x^2 + 1) + C$$

5. Solve the initial value problem if  $f'(x) = x + e^x$  and  $f(0) = 4$ .

6. A particle moving along the  $x$ -axis has acceleration  $a(t) = 10 \sin(t)$  measured in  $cm/s^2$ . Assume the particle as initial velocity  $v(0) = 0$  and initial position  $s(0) = 0$ , find a function that models its velocity,  $v(t)$ , and its position  $s(t)$ .